



1.1x Pre-Aliquoted Extensor Hi-Fidelity PCR Master Mix, 25µl Reaction

Description: Extensor Hi-Fidelity PCR Master Mix is a pre-aliquoted ready-to-use enzyme mix for long and accurate PCR without the need to mix individual components, reducing the risk of contamination and pipetting errors. The Extensor PCR Enzyme Mix, dNTPs, Extensor Reaction Buffer and MgCl₂ are all present in the mix.

Ordering Information: Select the buffer type from Table 1 and consumable type and size from Table 2.

Cat. No.	Buffer Type
EX1-xxx-2	Buffer 1 - For templates <12kb
EX2-xxx-2	Buffer 2 - For templates >12kb or problematic amplifications of any length.

Note: xxx denotes plastic consumable type

Cat. No.	Description	Quantity
EX?-266-2/a	0.2ml Thermo-Strips & Domed Caps	25µl PCR x 12 strips of 8 tubes
EX?-266-2/b	0.2ml Thermo-Strips & Domed Caps	25µl PCR x 10 x 12 strips of 8 tubes
EX?-337-2/a	Individual 0.2ml Domed Cap Thermo-Tubes	25µl x 96 tubes
EX?-337-2/b	Individual 0.2ml Domed Cap Thermo-Tubes	25µl x 10 x 96 tubes
EX?-350-2/a	Individual 0.5ml Flat Cap Thermo-Tubes	25µl x 96 tubes
EX?-350-2/b	Individual 0.5ml Flat Cap Thermo-Tubes	25µl x 10 x 96 tubes
EX?-384-2	Thermo-Fast [®] 384 Plates	25µl x 10 plates of 384 wells
EX?-489-2/a	Individual 0.5ml Domed Cap Thermo-Tubes	25µl x 96 tubes
EX?-489-2/b	Individual 0.5ml Domed Cap Thermo-Tubes	25µl x 10 x 96 tubes
EX?-600-2	0.2ml Thermo-Fast [®] 96 Plates	25µl x 10 plates of 96 wells
EX?-620-2/a	Individual 0.2ml Flat Cap Thermo-Tubes	25µl x 96 tubes
EX?-620-2/b	Individual 0.2ml Flat Cap Thermo-Tubes	25µl x 10 x 96 tubes
EX?-700-2	0.2ml Low Profile Thermo-Fast [®] 96 Plates	25µl x 10 plates of 96 wells
EX?-772-2/a	0.2ml Low Profile Thermo-Strips & Domed Caps	25µl PCR x 12 strips of 8 tubes
EX?-772-2/b	0.2ml Low Profile Thermo-Strips & Domed Caps	25µl PCR x 10 x 12 strips of 8 tubes
EX?-773-2/a	0.2ml Low Profile Thermo-Strips & Flat Caps	25µl PCR x 12 strips of 8 tubes
EX?-773-2/b	0.2ml Low Profile Thermo-Strips & Flat Caps	25µl PCR x 10 x 12 strips of 8 tubes
EX?-800-2	0.2ml Skirted Thermo-Fast [®] 96 Plates	25µl x 10 plates of 96 wells
EX?-848-2/a	0.2ml Low Profile Thermo-Strips & Domed Caps	25µl PCR x 8 strips of 12 tubes
EX?-848-2/b	0.2ml Low Profile Thermo-Strips & Domed Caps	25µl PCR x 10 x 8 strips of 12 tubes
EX?-849-2/a	0.2ml Low Profile Thermo-Strips & Flat Caps	25µl PCR x 8 strips of 12 tubes
EX?-849-2/b	0.2ml Low Profile Thermo-Strips & Flat Caps	25µl PCR x 10 x 8 strips of 12 tubes

For research purposes only



Cat. No.	Description	Quantity
EX?-900-2	0.2ml Semi Skirted Thermo-Fast® 96 Plates	25µl x 10 plates of 96 wells
EX?-937-2	Thermo-Fast® 384 Mark II Plates	25µl x 10 plates of 384 wells
EX?-990-2	0.2ml Ultra Rigid Semi Skirted Thermo-Fast® 96 Plates	25µl x 10 plates of 96 wells
EX?-1000-2	0.2ml Ultra Rigid Skirted Thermo-Fast® 96 Plates	25µl x 10 plates of 96 wells
EX?-1100-2	0.2ml Thermo-Fast® 96 PCR Detection Plate	25µl x 10 plates of 96 wells
EX?-1300-2	Thermo-Fast® 96 Robotic PCR Plate, Capped	25µl x 10 plates of 96 wells
EX?-1400-2	Thermo-Fast® 96 PCR Detection Plate Mark II, Capped	25µl x 10 plates of 96 wells
EX?-1111-2	Thermo-Fast® 384 Diamond PCR Plate	25µl x 10 plates of 384 wells
EX?-1113-2/a	0.2ml Thermo-Strips with Domed Caps	25µl PCR x 8 strips of 12 tubes
EX?-1113-2/b	0.2ml Thermo-Strips with Domed Caps	25µl PCR x 10 x 8 strips of 12 tubes
EX?-1114-2/a	0.2ml Thermo-Strips with Flat Caps	25µl PCR x 8 strips of 12 tubes
EX?-1114-2/b	0.2ml Thermo-Strips with Flat Caps	25µl PCR x 10 x 8 strips of 12 tubes

Note: Replace ? with buffer type

Example: **EX1-337-2/a**
1.1x Extensor Hi-Fidelity PCR Master Mix with Buffer 1, (Reaction Volume 25µl) Pre-aliquoted into 96 x 0.2ml Domed Cap Thermo-Tubes

Kit Components: Each well contains 22.5µl of a 1.1x working concentration PCR Master Mix. The addition of the template and primers (in a volume of 2.5µl) results in a final reaction volume of 25µl, containing:

Buffer 1

1x Extensor Buffer 1
 1.25 units DNA Polymerase
 2.25mM MgCl₂
 350µM each of dATP, dCTP, dGTP and dTTP

Buffer 2

1x Extensor Buffer 2
 1.25 units DNA Polymerase
 2.25mM MgCl₂
 500µM each of dATP, dCTP, dGTP and dTTP

Thermo-Fast® 96 plates are provided capped with Domed Cap Strips. An extra set of caps for application after the addition of template and primers is included with both plates and Thermo-Strips. Thermo-Fast® 384 plates are supplied heat sealed with Easy Peel (or Diamond Foil Seal for Diamond plates).

For research purposes only



Overview of Protocol:

For a 25µl reaction, take 22.5µl of Extensor Master Mix and add template, primers and water in a 2.5µl volume (scale up or down accordingly if required). Generally, 100–250ng template DNA, and 200nM (final concentration) of each primer is added. It is recommended that the Extensor Master Mix and added components are kept on ice. This removes the need for a hot start, as well as avoiding any degradation of primers and template through the 3' to 5' proofreading activity present in the Extensor Master Mix. The use of wax is not recommended, as it prevents adequate mixing of reaction components, leading to low yields. The enzyme mix has a fidelity that is at least four times higher than standard *Taq* DNA polymerase. All reaction tubes should be sterile and certified DNase/RNase free. The following points should also be noted:

- The Extensor Hi-Fidelity PCR Master Mix offers very robust amplification up to 15kb of human genomic DNA. Above 15kb, more optimisation may be required.
- Ensure proper mixing of reaction components, and always use thin-walled PCR tubes.
- Use a mineral oil overlay unless a heated lid thermocycler is used.
- Touchdown PCR may increase PCR product specificity.
- For best results, use primers (lengths 22–34 nucleotides) with annealing temperatures over 60°C.
- Primers can be used at 400nM for very long extensions.

Templates:

For the amplification of large DNA fragments, the quality of the template DNA is very important, as are the denaturation conditions. Keep template DNA denaturation steps as short as possible. Use Extensor Buffer 2 for DNA templates ≥ 12kb and when difficulties are expected or encountered. 125ng human genomic DNA is generally sufficient to provide good PCR results. When using simple templates (such as λ DNA), 1–10ng template DNA should prove sufficient; the number of cycles may be reduced by 5 and Extensor Buffer 1 can be used.

For research purposes only



Thermal Cycler Programming:

For high fidelity PCR, a standard protocol should be used. For long PCR, modifications may need to be made. An example of a long PCR thermal cycling programme is given:

Initial denaturation	92-94°C ¹	2 min	1 cycle
Denaturation	92-94°C	10 sec	
Annealing ²	50-68°C	30 sec	10 cycles
Extension	68°C ³	x min ⁴	
Denaturation	94°C	20 sec	
Annealing ²	50-68°C	30 sec	15-20 cycles
Extension	68°C	x min (+20 s/cycle)	
Final extension	68°C	7 min	1 cycle

1 - When amplifying over 15kb, use a denaturation temperature of 92°C.

2 - Annealing temperature dependent on primers.

3 - Always use an extension temperature of 68°C, if possible. Often good results are obtained using a single annealing/extension step at 68°C.

4 - Extension times depend on the length of sequence to be amplified (see table below).

Amplicon size (kb)	3	6	10	20	30	40
Extension time (min)	2	4	8	15	20	30

After PCR, a sample (10–30% of reaction) may be loaded directly on a gel.

Troubleshooting: **1 No product detected**

Try reducing the annealing temperature, increasing the concentration or quality of template, number of cycles or improving the purity of primers used.

2 Spurious bands appearing on electrophoresis gel

When non-specific products are amplified, try increasing the annealing temperature (up to a maximum of 68°C) or reducing template concentration or cycle number.

For research purposes only



Storage Conditions: Can be stored at -20°C in a constant temperature freezer for up to 1 year. Avoid freeze thawing. Once opened, the vial can be stored at 4°C for up to 1 month. Shipped on ice.

A license under the foreign counterparts of U.S. Patents Nos. 4,683,202, 4,683,195 and 4,965,188, owned by F. Hoffman-La Roche Ltd, for use in research and development, has an up-front fee component and a running royalty component. The purchase price of this product includes limited, non-transferable rights under the running-royalty component to use only this amount of the product to practice the Polymerase Chain Reaction ("PCR") and related processes described in said patents solely for the research and development activities of the purchaser when this product is used in conjunction with a thermal cycler whose use is covered by the up-front fee component. Rights to the up-front fee component must be obtained by the end user in order to have a complete license. These rights under the up-front fee component may be purchased from Perkin-Elmer or obtained by purchasing an Authorized Thermal Cycler. No right to perform or offer commercial services of any kind using PCR, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is hereby granted by implication or estoppel. Further information on purchasing licenses to practice the PCR process may be obtained by contacting the Director of Licensing at the Perkin-Elmer Corporation, 850 Lincoln Centre Drive, Foster City, California 94404

For research purposes only